

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry area away from heat or open flame and inaccessible to children.

PESTICIDE DISPOSAL: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Do not puncture or incinerate! **If empty:** Place in trash or offer for recycling if available. **If partly full:** Call your local solid waste agency for disposal instructions.

Contains no CFCs or other ozone depleting substances. Federal regulations prohibit CFC propellants in aerosols.



CONDITIONS OF SALE AND WARRANTY

Follow the **Directions for Use**. It is impossible to eliminate all risks inherently associated with use of this product, and therefore all such risk shall be assumed by the Buyer. Whitmire warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions for Use**, subject to the inherent risks, referred to above. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW: (A) WHITMIRE MAKES NO OTHER WARRANTIES EXPRESS OR IMPLIED, INCLUDING WARRANTIES OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY, (B) BUYER'S EXCLUSIVE REMEDY AND WHITMIRE'S AND SELLER'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT, AND (C) WHITMIRE AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, INCIDENTAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. Whitmire and the Seller offer this product, and the Buyer accepts it, subject to these **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of Whitmire.

Manufactured for:

Whitmire Micro-Gen Research
Laboratories, Inc.®

by BASF Corporation

3568 Tree Court Industrial Blvd.

St. Louis MO 63122-6682

Questions? 1-800-777-8570

© 2011 Whitmire Micro-Gen Research
Laboratories, Inc.®

81056237

110119-11

305xxxxx

Prescription Treatment® brand

Alpine®

Flea Insecticide with IGR

KILLS: Adult & Immature fleas. Kills fleas for up to 30 days. Kills hatching flea eggs for up to 7 months.

FOR USE IN: Apartments, Commercial Structures, Homes, Hotels, Kennels, Motels and Veterinary Clinics

ACTIVE INGREDIENTS:

Dinotefuran: N-methyl-N-
[(tetrahydro-3-furanyl)methyl]guanidine 0.25%
Pyriproxyfen 0.10%
Prallethrin 0.05%

OTHER INGREDIENTS: 99.60%
TOTAL: 100.00%

EPA Reg. No. 499-540 • EPA Est. No. 7969-MO-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

NET CONTENTS: 1 lb, 4 oz (20 oz)

FIRST AID

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 min. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN: May pose an aspiration pneumonia hazard. Contains petroleum distillate.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call 1-800-832-HELP for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Contains Petroleum Distillate. Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

See side panel(s) for Directions for Use
and Storage and Disposal.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

DIRECTIONS FOR USE

**IT IS A VIOLATION OF FEDERAL LAW
TO USE THIS PRODUCT IN A MANNER
INCONSISTENT WITH ITS LABELING.**

PRODUCT INFORMATION

This product kills fleas and prevents flea infestations for a full season. Adult fleas may be seen in the treated areas when brought in on infested animals or when adults emerge from pupal cases; however, populations will not develop in the treated area.

In transportation equipment, certain plastics may be whitened or dulled. Apply only to carpets and cloth upholstered areas, or test an inconspicuous area before treating.

NOTICE: This product has been tested on several types of carpet, fabric and other household furnishings without adverse affects. In a few instances, waxed surfaces have been whitened or dulled. Holding the can 36" away from the target surface normally prevents any whitening from occurring. Because of the wide variety of floor types and finishes, treat a small inconspicuous area before treating the entire area. Avoid contact with antique finishes.

APPLICATION INSTRUCTIONS

SHAKE WELL BEFORE USE! Use at the rate of 20 oz [one can] for up to 2,625 ft² (10 oz [half can] for up to 1,300 ft²). Vacuum before treatment and after treatment has dried. Treat infested areas, or potentially infested areas, including rugs, carpets, upholstered furniture, pet beds and pet resting areas. Hold can at arm's length and direct spray toward the area to be treated. Use a sweeping motion to apply product and back away from treated area while holding the can 36" away from the surface being treated. An area of 80 – 100 ft² can be treated in approx. 10 sec with this product. When treating upholstered furniture, treat under cushions and areas where flea development occurs.

USER RESTRICTIONS

- Cover aquariums and fish bowls and remove birds from area prior to treating.
- Vacate areas to be treated and do not reoccupy or contact treated surfaces until dry.
- Do not apply in food/feed areas.

TO CONTROL FLEAS IN PET AREAS: Remove old bedding and replace with fresh, clean bedding after treatment. **Do not treat pets with this product.** Treat pets with a product registered for use for flea control on animals in conjunction with this treatment.